

Four new species of the jumping spider genus *Portia* (Araneae, Salticidae) from China

Xin Xu¹, Xianjin Peng¹, Daiqin Li²

1 College of Life Sciences, Hunan Normal University, Changsha 410081, Hunan, China **2** Department of Biological Sciences, National University of Singapore, Singapore 117543, Singapore

Corresponding authors: Xin Xu (xuxin@hunnu.edu.cn), Daiqin Li (dbslidq@nus.edu.sg)

Academic editor: Jeremy Miller | Received 10 August 2021 | Accepted 5 October 2021 | Published 3 November 2021

<http://zoobank.org/7003C83E-5FB9-423F-82FB-F83DAF7C976D>

Citation: Xu X, Peng X, Li D (2021) Four new species of the jumping spider genus *Portia* (Araneae, Salticidae) from China. ZooKeys 1068: 27–40. <https://doi.org/10.3897/zookeys.1068.72805>

Abstract

We diagnose and describe four new species of *Portia* Karsch, 1878 and describe for the first time the male of *P. zhaoi* Peng, Li & Chen, 2003 from China based on morphological characters. The females of *Portia bawang* sp. nov. have the narrowest epigyne orifice. The males of *Portia erlangping* sp. nov. have the shortest embolus among all the species. The females of *Portia fajing* sp. nov. can be distinguished from other species by the anterior orifice margin, which is nearly parallel to the posterior orifice margin. The males of *Portia xishan* sp. nov. can be identified by the tegular furrow which extends to form a membrane and by the tegular apophysis which is obscured; the females of *Portia xishan* sp. nov. can be diagnosed by the slit-like epigynal orifice. The males of *P. zhaoi* have the longest embolus among all the species, and females can be diagnosed by the circular epigyne orifice and the longest copulatory ducts. To facilitate future identification, we also provide the GenBank accession codes of the DNA barcode gene, Cytochrome c oxidase subunit I (COI), for all the type specimens.

Keywords

Description, morphology, Hainan, Hong Kong, taxonomy

Introduction

Portia Karsch, 1878 is the most thoroughly studied jumping spider genus and one of the best-known model systems for behavioural and evolutionary research in spiders (Su et al. 2007; Harland et al. 2012). Unlike typical jumping spiders, species of *Portia* are both

cursorial predators and web builders. They build large, three-dimensional prey-catch webs (Jackson 1985). They also prey on other spiders by invading their webs and using aggressive mimicry to trick, then catch the resident spider. In addition, *Portia* species also eat insects ensnared in the alien web. Furthermore, all species of *Portia* show specialized prey-catching behaviour for a particular type of prey and have a preference for spiders as prey over insects (reviewed by Jackson and Pollard 1996; Li and Jackson 1996; Harland et al. 2012).

Portia was erected based on the female morphology of *Portia schultzi* Karsch, 1878 (Karsch 1878). A taxonomic revision of the whole genus was completed by Wanless (1978). The monophyly of *Portia* is now strongly supported by both molecular and morphological data (Su et al. 2007; Maddison et al. 2014; Maddison 2015). *Portia* belongs to the subfamily Spartaecinae, tribe Spartaecini, subtribe Spartaecina (Maddison 2015). *Portia* is sister to *Cyrrba* Simon, 1986 and *Paracyrrba* Zabka & Kovac, 1996 (Su et al. 2007; Maddison 2014). To date, the genus contains 17 species worldwide, mainly distributed in the Oriental and Ethiopian regions, and specifically, 10 out of 17 *Portia* species occur in China (Wanless 1978; World Spider Catalog 2021). Peng and Li (2002) reported a taxonomic review of Chinese *Portia* species. The key to species of *Portia* was provided in the studies of Wanless (1978) and Peng and Li (2002) based on male and female genital morphology. Since Peng et al. (2003), only one new species of *Portia* has been reported from Taiwan (Zhang and Li 2005; World Spider Catalog 2021). In this study, after examining the vouchers collected in China, we identify and describe four new species of *Portia* and describe the male of *P. zhaoi* for the first time based on male and/or female genital morphology.

Materials and methods

All specimens were collected from China (Fig. 1). We removed the right four legs of adults for molecular work, preserved them in 100% ethanol, and kept them at -80°C . We preserved the remains of each specimen in 80% ethanol as a voucher for morphological examination. All voucher specimens are deposited at the College of Life Sciences, Hubei University, Wuhan, Hubei Province, China.

We examined and dissected the specimens under an Olympus SZ51 stereomicroscope. The soft tissues of female genitalia were degraded using 10 mg/ml trypsin (Bomei Biotech Company, Hefei, Anhui, China) for at least 3 h at room temperature. Male and female genitalia were photographed with a digital camera CCD mounted on an Olympus BX53 compound microscope, and then generated compound focused images with Helicon Focus v. 6.7.1. All measurements were made using a digital camera MC170HD mounted on a Leica M205C stereomicroscope and are given in millimeters. Leg and palp measurements are given in the following order: leg total length (femur + patella + tibia + metatarsus + tarsus), palp total length (femur + patella + tibia + tarsus).

Abbreviations used: AL = abdomen length; ALE = anterior lateral eyes; AME = anterior median eyes; AW = abdomen width; BL = body length; CF = cymbium flange; CL = carapace length; CW = carapace width; E = embolus; PLE = posterior lateral eyes; PME = posterior median eyes; T = tegulum; TA: tegular apophysis; TF = tegular furrow.




Figure 1. Map showing the collection sites of five *Portia* species in this study.

Taxonomy

Genus *Portia* Karsch, 1878

Type species. *Portia schultzi* Karsch, 1878

Diagnosis. The genus *Portia* can be distinguished from other genera of the subfamily Spartaeinae by the dorsum of the abdomen with distinct tufts, the ventral tibiae with long fan-like fringes, and the malp palp with a dorsal cymbium flange (Zhang and Li 2005).

Portia bawang sp. nov.

<http://zoobank.org/3833BB68-349D-4268-9D0B-F977EDB1DFA3>

Figure 2

Type material. Holotype: CHINA • 1 ♀; Hainan Province, Changjiang County, Bawang National Forest Park; 19.023°N, 109.103°E, alt. 692 m; 19 July 2012; F.X. Liu, D. Li and X. Xu leg.; DL-002-013-2012. **Paratypes:** CHINA • 1 ♀; same data as for the holotype; 19.027°N, 109.101°E, alt. 702 m; 7 August 2017; F.X. Liu, D. Li and X. Xu leg.; DL-003-002-2017 • 2 ♀♀; Hainan Province, Ledong County, Jianfeng National




Figure 2. General somatic morphology and female genital anatomy of *Portia bawang* sp. nov. **A–D** DL-002-013-2012 (holotype) **A, B** female **C** epigyne, ventral view **D** vulva, dorsal view. Scale bar: 0.3 mm.

Forest Park; 19.296°N, 109.074°E, alt. 565–1005 m; 22 July 2012; F.X. Liu, D. Li and X. Xu leg.; DL-002-018-2012, DL-002-022-2012 • 1 ♀; Hainan Province, Lingshui County, Diaoluo National Forest Park; 18.400°N, 109.559°E, alt. 105 m; 21 June 2011; D. Li leg.; DL-005-006-2011.

Diagnosis. Females of *P. bawang* sp. nov. resemble those of *P. fimbriata*, *P. quei*, and *P. taiwanica* but can be distinguished from them by the epigyne orifice being narrowest (Fig. 2C; for comparison with known species, see fig. 8 in Wanless (1978), figs 664 and 666 in Peng et al. (1993), and fig. 4F in Zhang and Li (2005), respectively); from those of *P. labiata* by the slightly straight anterior margin of posterior depression (Fig. 2C, D; see fig. 1C in Zhu et al. (2007)); from those of *P. heteroidea* by lacking a median septum (Fig. 2C; see figs 10–12 in Xie and Yin (1991)); from those of *P. fajing* sp. nov. and *P. xishan* sp. nov. by the epigyne orifice being narrowest and elliptical (Fig. 2C); from those of *P. zhaoi* by a smaller elliptical epigyne orifice and a shorter copulatory duct (Fig. 2C, D).

Description. Female (holotype; Fig. 2A, B). Carapace greyish brown; ocular area yellow brown, with tufts of orange brown hairs around AME. Clypeus brown with dense ventral white hairs. Chelicerae dark brown with 3 small promarginal and 3 large retro-

marginal teeth. Maxillae and labium black-brown with reddish brown anterior margin. Sternum light brown, densely covered with creamy white hairs. Measurements: eye sizes: AME 0.80, ALE 0.31, PME 0.23, PLE 0.32, anterior eye row 2.46 wide, posterior eye row 2.30 wide, eye area 1.70 long; clypeus height 0.57; BL 8.17–9.46; holotype BL 9.46, CL 4.19, CW 3.09, AL 5.55, AW 3.68; palp 3.96 (1.27 + 0.58 + 0.76 + 1.35), leg I 12.56 (3.19 + 1.57 + 3.06 + 3.22 + 1.52), leg II 10.25 (2.95 + 1.41 + 2.26 + 2.41 + 1.22), leg III 8.48 (2.40 + 1.20 + 1.67 + 2.11 + 1.10), leg IV 13.89 (3.59 + 1.05 + 3.15 + 4.74 + 1.36). Leg formula 4123. Legs slender, ventral portion of tibiae fringed with long black hairs. Dorsum of abdomen black brown, anterior portion light brown with grey-white hairs, middle portion with a small patch and posterior portion with two oval patches, the three patches densely covered with greyish long hairs.

Female genitalia. Epigyne orifice undivided, highly sclerotised, elliptical, anterior orifice margin distinct and posterior margin wide, slightly curved; spermathecae large and spherical (Fig. 2C, D).

Etymology. The species epithet, a noun in apposition, refers to the type locality.

Distribution. Hainan (Changjiang, Ledong, Lingshui).

GenBank accession code of holotype. OK235444.

***Portia erlangping* sp. nov.**

<http://zoobank.org/8492FA45-A5CB-4000-95EB-6CD65D54C523>

Figure 3

Type material. Holotype: CHINA • 1 ♂; Henan Province, Nanyang City, Xixia County, Erlangping Town; 33.524°N, 111.688°E; 11 April 2013; F.X. Liu leg.; HN-013-001.

Paratype: CHINA • 1 ♂; same data as for the holotype; HN-013-002.

Diagnosis. Males of *P. erlangping* sp. nov. resemble those of *P. heteroidea* but can be distinguished from the latter by the tegulum having one curved furrow (Fig. 3A), while *P. heteroidea* has two furrows (see fig. 6 in Xie and Yin (1991)), and by the longer cymbium flange (Fig. 3D; see fig. 8 in Xie and Yin (1991)); from those of *P. albimana* by the longer embolus, larger cymbium flange and thicker retrolateral tibial apophysis (Fig. 3A, D; see fig. 12B–D in Wanless (1978)); from those of *P. assamensis*, *P. fimbriata*, *P. labiata*, *P. orientalis*, *P. quei*, *P. xishan* sp. nov., *P. taiwanica*, and *P. zhaoi* by the embolus being shortest (Fig. 3A; see figs 10D, 7C, and 10A in Wanless (1978), fig. 6 in Murphy and Murphy 1983, fig. 661 in Peng et al. (1993), and fig. 4B in Zhang and Li (2005), respectively); in addition, from those of *P. assamensis* and *P. fimbriata* by the cymbium flange being thickest and longest (Fig. 3D; figs 10E, 7G in Wanless (1978), respectively); from those of *P. labiata* by the larger cymbium flange and thicker retrolateral tibial apophysis (Fig. 3D; see fig 10B in Wanless (1978)); from those of *P. orientalis*, *P. quei*, *P. taiwanica* and *P. zhaoi* by the bar-shaped retrolateral tibial apophysis (Fig. 3D; see fig. 6 in Murphy and Murphy 1983, and fig. 4D in Zhang and Li (2005), respectively).

Description. Male (holotype). Carapace greyish brown with white band on thoracic groove and lateral margin. Ocular area yellow-brown, with tufts of yellow-brown




Figure 3. Male genital anatomy of *Portia erlangping* sp. nov. **A–D** HN-013-001 (holotype) **A** palp, ventral view **B, C** palp, retrolateral view **D** palp, dorsal view. Scale bars: 0.3 mm.

hairs around AME. Clypeus black-brown without dense ventral white hairs. Chelicerae dark brown with 2 small promarginal and 3 large retromarginal teeth. Maxillae and labium black-brown with white anterior margin. Sternum black-brown, densely covered with creamy white hairs. Measurements: eye sizes: AME 0.64, ALE 0.24, PME 0.17, PLE 0.21, anterior eye row 1.78 wide, posterior eye row 1.65 wide, eye area 1.38 long; clypeus height 0.30; BL 5.84–6.45; holotype BL 5.84, CL 2.89, CW 2.14, AL 2.95, AW 1.60; leg I 8.40 (2.25 + 0.92 + 2.02 + 2.11 + 1.10), leg II 6.95 (1.90 + 0.94 + 1.57 + 1.60 + 0.94), leg III 6.46 (1.92 + 0.74 + 1.30 + 1.72 + 0.78), leg IV 9.45 (2.47 + 0.97 + 2.06 + 2.94 + 1.01). Leg formula 4123. Legs black-brown, slender, ventral portion of tibiae fringed with long black hairs. Dorsum of abdomen black-brown, anterior portion light brown with grey-white hairs, middle portion with a small triangular patch and posterior portion with two oval patches, the three patches densely covered with grey-white hairs.

Palp. Tibia with 3 apophyses, ventral one thick and short, intermediate one relatively slender, retrolateral one largest and bar-shaped in dorsal view (Fig. 3A–D). Embolus short and stout (Fig. 3A). Seminal duct clear and S-shaped. Tegulum with a deeply curved furrow and a membranous apophysis (Fig. 3A, B). Cymbium flange robust, terminal portion overlapping on base of retrolateral tibial apophysis dorsally (Fig. 3D).

Etymology. The species epithet, a noun in apposition, refers to the type locality.

Distribution. Henan (Nanyang)

Portia fajing sp. nov.

<http://zoobank.org/E903C81D-1F7B-4966-91DD-93EB51627C4A>

Figure 4

Type material. Holotype: CHINA • 1 ♀; Zhejiang Province, Hangzhou City, Fajing Temple; 30.234°N, 120.095°E; alt. 79 m; 13 July 2013; F.X. Liu, D. Li, X. Xu and Z.T. Zhang leg.; DL-001-016-2013. **Paratypes:** CHINA • 4 ♀♀; same data as for the



Figure 4. Female genital anatomy of *Portia fajing* sp. nov. **A, B** DL-001-016-2013 (holotype) **A** epigyne, ventral view **B** vulva dorsal view. Scale bar: 0.3 mm.

holotype, 30.228°N, 120.091°E; alt. 117 m; DL-001-015-2013, DL-002-016-2013, DL-003-016-2013, DL-003-017-2013.

Diagnosis. Females of *P. fajing* sp. nov. can be distinguished from those of *P. bawang* sp. nov., *P. labiata*, *P. quei*, and *P. taiwanica* by the slit-like epigynal orifice, the anterior orifice margin nearly parallel to the posterior orifice margin ventrally, and the W-shaped posterior epigynal margin (Fig. 4A, B; see fig. 1C in Zhu et al. (2007), fig. 664 and 665 in Peng et al. (1993), and fig. 4F in Zhang and Li (2005), respectively); from those of *P. xishan* sp. nov. by the W-shaped posterior orifice margin (Fig. 4B); from those of *P. fimbriata* and *P. zhaoi* by the copulatory duct being shortest (Fig. 4B; see fig. 8D, E in Wanless (1978))

Description. Female (holotype). Carapace brown; ocular area yellow-brown, with tufts of orange-brown hairs around AME. Clypeus brown with dense ventral white hairs. Chelicerae dark brown with 3 small promarginal and 3 large retro-marginal teeth. Maxillae and labium reddish brown with yellow-brown anterior margin. Sternum yellow-brown, densely covered with creamy white hairs. Measurements: eye sizes: AME 0.68, ALE 0.30, PME 0.21, PLE 0.29, anterior eye row 2.01 wide, posterior eye row 1.87 wide, eye area 1.55 long; clypeus height 0.35; BL 6.56–7.64; holotype BL 7.64, CL 3.23, CW 2.58, AL 4.71, AW 3.22; palp 2.80 (0.77 + 0.48 + 0.59 + 0.96), leg I 8.74 (2.37 + 1.10 + 2.01 + 2.23 + 1.03), leg II 7.17 (2.01 + 1.06 + 1.60 + 1.61 + 0.89), leg III 6.61 (1.83 + 0.90 + 1.35 + 1.74 + 0.79), leg IV 10.84 (2.76 + 1.12 + 2.43 + 3.62 + 0.91). Leg formula 4123. Legs brown, the ventral portion of tibiae fringed with long black hairs. Dorsum of abdomen greyish brown, posterior portion with two circular patches densely covered with grey-white long hairs.

Female genitalia. Epigyne orifice undivided, highly sclerotised, transverse, slit-like, posterior orifice margin W-shaped; spermatheca large and spherical (Fig. 4B).

Etymology. The species epithet, a noun in apposition, refers to the type locality.

Distribution. Zhejiang (Hangzhou).

GenBank accession code of holotype. OK235443.

***Portia xishan* sp. nov.**

<http://zoobank.org/8B8AD618-B4D9-4128-82D2-033436119365>

Figure 5

Type material. Holotype: CHINA • 1 ♂; Yunnan Province, Kunming City, Western Mountains; 24.962°N, 102.631°E, alt. 2172 m; 16 August 2006; F.X. Liu and Q.Q. Liu leg.; LQ-18-06. **Paratypes:** CHINA • 1 ♂ 3 ♀♀; same data as for the holotype; LQ-18-06A/06B/06C/06D • 8 ♀♀; same data as for the holotype; 11 November 2020; L. Yu and X.R. Miao leg.; P2020001, P2020010, P2020011, P2020034, P2020039, P2020053, P2020054, P2020055.

Diagnosis. Males of *P. xishan* sp. nov. can be distinguished from those of *P. albimana* by the longer embolus, larger cymbium flange and thicker, bar-shaped retrolateral tibial apophysis (Fig. 5A–D; see fig. 12B–D in Wanless (1978)); from those of *P. assamensis*, *P. erlangping* sp. nov., *P. fimbriata*, *P. labiata*, *P. orientalis*, *P. quei*, *P. taiwanica*, and *P. zhaoi* by the tegular furrow extending a membranous apophysis ventrally, the tegular apophysis obscure, and the embolus basally with a spinule (Fig. 5A–C; see figs 10D, 7C, and 10A in Wanless (1978), fig. 6 in Murphy and Murphy 1983, fig. 661 in Peng et al. (1993), and fig. 4B in Zhang and Li (2005), respectively); in addition, from those of *P. assamensis* and *P. fimbriata* by the embolus being shortest (Fig. 5A–C; see figs 10D and 7C in Wanless (1978)); from those of *P. erlangping* sp. nov. by the embolus being longer (Fig. 5A); from those of *P. labiata* by the shorter cymbium flange and thicker bar-shaped retrolateral tibial apophysis (Fig. 4D; see fig. 10B in Wanless (1978)); from those of *P. orientalis*, *P. quei*, *P. taiwanica*, and *P. zhaoi* by the longer embolus and bar-shaped retrolateral tibial apophysis (Fig. 5A–D; see fig. 6 in Murphy and Murphy 1983, fig. 661 in Peng et al. (1993), and fig. 4B in Zhang and Li (2005), respectively). Females of *P. xishan* sp. nov. differ from those of *P. bawang* sp. nov. by having a slightly wider epigynal orifice (Fig. 5E); from those of *P. fajing* sp. nov. by the larger epigyne orifice and distinctly sclerotised anterior orifice margin (Fig. 5E); from those of *P. labiata*, *P. quei*, and *P. taiwanica* by the slit-like epigynal orifice (Fig. 5E; see fig. 1C in Zhu et al. (2007), fig. 664 and 665 in Peng et al. (1993), and fig. 4F in Zhang and Li (2005), respectively); from those of *P. fimbriata* and *P. zhaoi* by the copulatory duct being shortest (Fig. 5F; see fig. 8D, E in Wanless (1978)).

Description. Male (Holotype). Carapace black-brown with white band on thoracic groove and lateral margin. Ocular area yellow-brown, with tufts of yellow-brown hairs around AME. Clypeus black-brown without dense ventral white hairs. Chelicerae dark brown with 5 small promarginal and 2 large retromarginal teeth. Maxillae and labium black-brown with yellow-brown anterior margin. Sternum yellow-brown, densely covered with creamy white hairs. Measurements: eye sizes: AME 0.59, ALE 0.30, PME 0.19, PLE 0.28, anterior eye row 1.78 wide, posterior eye row 1.69 wide, eye area 1.32 long; clypeus height 0.42; BL 5.72–6.21; holotype BL 5.72, CL 2.87, CW 2.17, AL 2.75, AW 1.54; leg I 8.02 (2.04 + 0.91 + 1.81 + 2.04 + 1.22), leg II 6.80 (1.95 + 0.81 + 1.38 + 1.71 + 0.95), leg III 6.58 (1.87 + 0.86 + 1.29 + 1.64 + 0.92),




Figure 5. Male and female genital anatomy of *Portia xishan* sp. nov. **A–D** LQ-18-06 (holotype) **A** palp, ventral view **B, C** palp, retrolateral view **D** palp, dorsal view **E, F** LQ-18-06B **E** epigyne, ventral view **F** vulva, dorsal view. Scale bars: 0.3 mm.

leg IV 9.85 (2.61 + 0.92 + 2.09 + 3.17 + 1.06). Leg formula 4123. Legs black-brown, slender, the ventral portion of tibiae fringed with long black hairs. Dorsum of abdomen greyish brown, anterior portion light brown with grey-white hairs, three pairs of oval patches densely covered with grey-white hairs, the posterior pair largest.

Palp. Tibia with 3 apophyses, ventral one thick and short, intermediate one relatively slender, retrolateral one largest and bar-shaped in dorsal view (Fig. 5A–D). Embolus short and stout, with a spinule basally in retrolateral view (Fig. 5A–C). Seminal duct clear and S-shaped. Tegulum with a curved furrow extending a membranous apophysis ventrally and an obscure tegular apophysis (Fig. 5A–C). Cymbium flange robust, terminal portion close to middle portion of retrolateral tibial apophysis dorsally (Fig. 5D).

Female (LQ-18-06B). Carapace yellow-brown; ocular area yellow-brown, with tufts of greyish brown hairs around AME. Clypeus brown with densely ventral white hairs. Chelicerae dark brown with 4 small promarginal and 3 large retromarginal teeth. Maxillae and labium black-brown with yellow brown to white hairs on anterior margin. Sternum brown, densely covered with creamy white hairs. Meas-

measurements: eye sizes: AME 0.67, ALE 0.30, PME 0.23, PLE 0.27, anterior eye row 1.93 wide, posterior eye row 1.86 wide, eye area 1.67 long; Clypeus height 0.42; BL 5.83–7.66; LQ-18-06B: BL 6.10, CL 2.79, CW 2.44, AL 3.42, AW 2.11; palp 2.93 (0.95 + 0.31 + 0.58 + 1.09), leg I 7.20 (2.09 + 0.88 + 1.70 + 1.52 + 1.01), leg II 6.17 (2.01 + 0.95 + 1.15 + 1.15 + 0.91), leg III 6.02 (1.82 + 0.72 + 1.19 + 1.39 + 0.90), leg IV 7.72 (2.85 + 0.78 + 1.46 + 1.71 + 0.92). Leg formula 4123. Legs brown, ventral portion of tibiae fringed with long black hairs. Dorsum of abdomen brown, anterior margin with numerous long white hairs, posterior portion with two circular patches densely covered with grey-white hairs.

Female genitalia. Epigyne orifice undivided, highly sclerotised, transverse, spindly, posterior orifice margin slightly curved; spermathecae large and spherical (Fig. 5E, F).

Etymology. The species epithet, a noun in apposition, “xishan” means Western Mountains in Chinese and refers to the type locality.

Distribution. Yunnan (Kunming).

GenBank accession code of holotype. OK235446.

Portia zhaoi Peng, Li & Chen, 2003

Figure 6

Portia zhaoi Peng, Li & Chen, 2003: 50, figs 1–4; Peng 2020: 356, fig. 255a–d.

Type material examined. Holotype: CHINA • 1 ♀; Guangxi Zhuang Autonomous region, Dongxing County, Rongguang Tea Plantation; 21.29°N, 108.02°E; 13 August 1992; F.X. Liu leg.

Additional material examined. CHINA • 1 ♂ 1 ♀; Hainan Province, Wuzhishan City, Shuiman Town, Yongxun Village; 18.903°N, 109.623°E, alt. 551 m; 25 July 2012; F.X. Liu, D. Li and X. Xu leg.; DL-002-024-2012, DL-001-024-2012; 1 ♂; Shenzhen, Xianhu Lake; 22.583°N, 114.169°E, alt. 66 m; 14 June 2012; F.X. Liu, D. Li and X. Xu leg.; DL-007-2012; • 1 ♀; Shenzhen, Yinhu Lake; 22.58°N, 114.08°E; 15 June 2012; F.X. Liu, D. Li and X. Xu leg.; DL-008-2012 • 1 ♂ 2 ♀♀; Hainan Province, Changjiang County, Bawang National Forest Park; 19.027°N, 109.101°E, alt. 702 m; 7 August 2017; F.X. Liu, D. Li and X. Xu leg.; LID-001-002-2017, LID-002-002-2017, LID-004-002-2017 • 1 ♀; Hong Kong, Kadoorie Farm and Botanic Garden; 22.424°N, 114.125°E, alt. 571 m; 12 April 2012; F.X. Liu, D. Li and X. Xu leg.; DL-001-2012 • 1 ♂; Hainan Province, Ledong County, Jianfeng Town, Institute of Tropical Forestry; 18.703°N, 108.789°E, alt. 129 m; 21 July 2012; F.X. Liu, D. Li and X. Xu leg.; DL-004-017-2012 • 1 ♂; Hainan Province, Yacha Town, 1st Burei Village; 19.193°N, 109.418°E, alt. 268 m; 18 July 2012; F.X. Liu, D. Li and X. Xu leg.; DL-011-2012.

Diagnosis. Males of *P. zhaoi* can be distinguished from those of all other *Portia* species by having the longest embolus (Fig. 6A); in addition, from those of *P. erlangping* sp. nov., *P. fimbriata* and *P. xishan* sp. nov. By the finger-shaped retrolateral tibial




Figure 6. Male and female genital anatomy of *Portia zhaoi* Peng, Li & Chen, 2003 **A–D** DL-002-024-2012 **A** palp, ventral view **B, C** palp, retrolateral view **D** palp, dorsal view **E, F** DL-001-024-2012 **G** holotype deposited in the Institute of Zoology, Chinese Academy of Sciences, Beijing **E, G** epigyne, ventral view **F** vulva dorsal view **H, I** male. Scale bars: 0.3 mm.

apophysis (Fig. 6A, B, D; see 7G in Wanless (1978)); from those of *P. quei* by the thicker ventral tibial apophysis (Fig. 6B; see fig. 661 in Peng et al. (1993)). Females of *P. zhaoi* can be distinguished from those of *P. labiata* by its longer copulatory ducts (Fig. 6E, F; see fig. 1D in Zhu et al. (2007)); from those of *P. bawang* sp. nov., *P. fajing* sp. nov., *P. fimbriata*, *P. quei*, *P. taiwanica*, and *P. xishan* sp. nov. by the circular epi-

gyne orifice (Fig. 6E; see fig. 8C–E in Wanless (1978), fig. 664 and 666 in Peng et al. (1993), and fig. 4F in Zhang and Li (2005), respectively).

Description. Male (DL-002-024-2012, Fig. 6G, H). Carapace yellow-brown with white band on thoracic groove and lateral margin. Ocular area yellow-brown, with tufts of yellow hairs around AME. Clypeus black-brown without dense ventral white hairs. Chelicerae dark brown with 4 small promarginal and 3 large retromarginal teeth. Maxillae and labium black-brown with yellow-brown anterior margin. Sternum yellow-brown, densely covered with creamy white hairs. Measurements: eye sizes: AME 0.68, ALE 0.29, PME 0.17, PLE 0.32, anterior eye row 1.88 wide, posterior eye row 1.67 wide, eye area 1.50 long; clypeus height 0.32; BL 5.62–7.63; holotype BL 5.69, CL 2.97, CW 2.32, AL 2.84, AW 1.32; leg I 9.24 (2.52 + 0.69 + 2.31 + 2.56 + 1.16), leg II 7.58 (2.26 + 0.86 + 1.61 + 2.02 + 0.83), leg III 6.54 (1.90 + 0.88 + 1.29 + 1.70 + 0.77), leg IV 9.54 (2.89 + 0.85 + 1.91 + 2.87 + 1.02). Leg formula 4123. Legs black brown, slender, the ventral portion of tibiae fringed with long black hairs. Dorsum of abdomen brown, anterior portion light brown with grey-white hairs, one pair of oval patches covered with dense grey-white hairs.

Palp. Tibia with 3 apophyses, ventral one thick and short, intermediate one relatively slender, retrolateral one largest and finger-shaped in dorsal view (Fig. 6A–D). Embolus slender and long (Fig. 6A). Seminal duct clear and S-shaped. Tegulum with a deeply curved furrow, and a triangular membranous apophysis (Fig. 6A–C). Cymbium flange robust, triangular, terminal portion overlapping on middle portion of retrolateral tibial apophysis dorsally (Fig. 6D).

Redescription. Female (DL-001-024-2012). Carapace black brown; ocular area yellow brown, with tufts of greyish hairs around AME. Clypeus brown with densely ventral white hairs. Chelicerae dark brown with 3 small promarginal and 3 large retromarginal teeth. Maxillae and labium black-brown with yellow brown anterior margin. Sternum yellow-brown, densely covered with creamy white hairs. Measurements: eye sizes: AME 0.79, ALE 0.35, PME 0.26, PLE 0.37, anterior eye row 2.29 wide, posterior eye row 2.13 wide, eye area 1.81 long; clypeus height 0.42; BL 7.18–8.24; DL-001-024-2012: BL 7.85, CL 4.05, CW 3.07, AL 4.17, AW 2.24; palp 3.64 (0.80 + 0.64 + 0.75 + 1.45), leg I 11.32 (3.16 + 1.39 + 2.64 + 2.92 + 1.21), leg II 9.52 (2.81 + 1.36 + 2.13 + 2.33 + 0.89), leg III 8.20 (2.39 + 1.05 + 1.67 + 2.23 + 0.86), leg IV 13.98 (3.71 + 1.44 + 2.98 + 4.94 + 0.91). Leg formula 4123. Legs black-brown, the ventral portion of tibiae fringed with long black hairs. Dorsum of abdomen brown, anterior margin with numerous long white hairs, posterior portion with three oval patches densely covered with grey-white hairs, the middle one small and the posterior two large.

Female genitalia. Epigyne orifice undivided, highly sclerotised, circular, anterior orifice margin distinct, posterior orifice margin slightly curved; spermathecae large and spherical (Fig. 6E, F).

Etymology. The species epithet, a noun in apposition, refers to the type locality.

Distribution. Guangxi, Hainan, Hong Kong, Shenzhen.

GenBank accession code of DL-002-024-2012. OK235445.

Acknowledgements

We thank Fengxiang Liu, Qingqing Liu, Long Yu, Gary Ades, Paul Crow, Yorkie Wong, Zoie Wong and Xinran Miao for their assistance in the field; Zhe Zhao for helping examine and photographing the female holotype of *P. zhaoi*; and Jeremy Miller, Suresh Benjamin, and two anonymous reviewers for their insightful comments on the manuscript. This study was supported by the National Natural Sciences Foundation of China (NSFC-32070430; NSFC-31872229), the Hunan Provincial Natural Science Foundation of China for Excellent Young Scholars (2021JJ20035) and the Singapore Ministry of Education AcRF Tier 1 grant (R-154-000-B72-114).

References

- Harland DP, Li D, Jackson RR (2012) How jumping spiders see the world. In: Lazareva O (Ed.) *How Animals See the World*. Oxford University Press, Oxford, 133–163. <https://doi.org/10.1093/acprof:oso/9780195334654.003.0010>
- Jackson RR (1985) Web-building jumping spiders. *Scientific American* 253: 102–115. <https://doi.org/10.1038/scientificamerican0985-102>
- Jackson RR, Pollard SD (1996) Predatory behavior of jumping spiders. *Annual Review of Entomology* 41: 287–308. <https://doi.org/10.1146/annurev.en.41.010196.001443>
- Karsch F (1878). Exotisch-araneologisches. *Zeitschrift für die Gesamten Naturwissenschaften* 51: 322–333, 771–826.
- Li D, Jackson RR (1996) Prey-specific capture behaviour and prey preferences of myrmicophagic and araneophagic jumping spiders (Araneae: Salticidae). In: Mahnert V (ed.) *Proceedings of the XIIIth International Congress of Arachnology*, Geneva, 3–8 September 1995. *Revue Suisse de Zoologie hors série* 2: 423–436.
- Maddison WP (2015) A phylogenetic classification of jumping spiders (Araneae: Salticidae). *Journal of Arachnology* 43: 231–292. <https://doi.org/10.1636/arac-43-03-231-292>
- Maddison WP, Li D, Bodner M, Zhang JX, Xu X, Liu QQ, Liu FX (2014) The deep phylogeny of jumping spiders (Araneae, Salticidae). *ZooKeys* 440: 57–87. <https://doi.org/10.3897/zookeys.440.7891>
- Murphy J, Murphy F (1983) More about *Portia* (Araneae: Salticidae). *Bulletin of the British Arachnological Society* 6: 37–45.
- Peng XJ, Li SQ (2002) Chinese species of the jumping spider genus *Portia* Karsch (Araneae: Salticidae). *Pan-Pacific Entomologist* 78: 255–264.
- Peng XJ, Li SQ, Chen J (2003) Description of *Portia zhaoi* sp. nov. from Guangxi, China (Araneae, Salticidae). *Acta Zootaxonomica Sinica* 28: 50–52.
- Peng XJ, Xie LP, Xiao XQ, Yin CM (1993) *Salticids in China* (Arachnida: Araneae). Hunan Normal University Press, Changsha, 270 pp.
- Su KFY, Meier R, Jackson RR, Harland DP, Li D (2007) Convergent evolution of eye ultrastructure and divergent evolution of vision-mediated predatory behavior in jumping

- spiders. *Journal of Evolutionary Biology* 20: 1478–1489. <https://doi.org/10.1111/j.1420-9101.2007.01335.x>
- Xie LP, Yin CM (1991) Two new species of Salticidae from China (Arachnida: Araneae). *Acta Zootaxonomica Sinica* 16: 30–34.
- Wanless FR (1978) A revision of the spider genus *Portia* (Araneae: Salticidae). *Bulletin of the British Museum of Natural History (Zoology)* 34: 83–124. <https://doi.org/10.5962/p.28739>
- World Spider Catalog (2021) World Spider Catalog. Version 22.5. Natural History Museum Bern. <http://wsc.nmbe.ch> [accessed on 2 August 2021]
- Zhang JX, Li D (2005) Four new and one newly recorded species of the jumping spiders (Araneae: Salticidae: Lyssomaninae & Spartaeinae) from (sub)tropical China. *Raffles Bulletin of Zoology* 53: 221–229. <http://doi.org/10.5281/zenodo.4619118>
- Zhu MS, Yang ZZ, Zhang ZS (2007) Description of a new species and a newly recorded species of the family Salticidae from China (Araneae: Salticidae). *Journal of Hebei University, Natural Science Edition* 27: 511–517.